

Plant Growth Regulators Impact on Vegetative and Reproductive Characteristics of Gladiolus Cut Flowers (*Gladiolus hybridus* Hort.)

Abdolali Hesami ^{1*} and Ali Dolatkahi ²

¹Persian Gulf University, Faculty of Agriculture and Natural Resources, Department of Horticultural Science, Bushehr, Iran

²Ferdowsi University of Mashhad, Faculty of Agriculture, Department of Horticultural Science, Mashhad, Iran

Received: 24 November 2015

Accepted: 30 January 2016

*Corresponding author's email: alihesami4400@yahoo.com

Gladiolus is an important marketable cut flower, especially valued for use in floral arrangements. We assessed the effects of two different methods of application (foliar spray and drench) and various concentrations of gibberellic acid (GA₃), salicylic acid (SA), mival (MI) and krizatcin (KR) on vegetative and reproductive features of *Gladiolus hybridus*. The experiment was planned in a factorial experiment based on a completely randomized design with four replicates. The results showed that plant growth regulators (PGRs) had significant effect on florets number, leaf area and corm weight and volume. The best treatment to increase the floral stalk length was 50 mg L⁻¹ GA₃ spray. Our results indicated that higher GA₃ doses significantly increased leaf area more than lower doses. Furthermore, maximum floret diameter was achieved at 100 mg L⁻¹ MI applied as foliar spray. The maximum number of florets (15.87) was obtained in the plants treated with intermediate concentrations of MI in both application methods. Corms drenched with 25 mg L⁻¹ MI effectively increased their corm weight and volume. Generally, MI was found to be more effective than other PGRs. Therefore, due to greater efficacy of mival, it is suggested to use this PGR in order to increase the growth of vegetative organs as well as reproductive attributes.

Abstract

Keywords: Application method, Gibberellin, Krizatcin, Mival, Salicylic acid.

Abbreviations: GA: Gibberellic acid, KR: Krizatcin, MI: Mival, PGRs: Plant growth regulators, SA: Salicylic acid.

INTRODUCTION

Gladiolus (Gladiolus hybridus Hort.) is one of the four famous cut flowers in the world, which is extensively cultivated in many countries (Ji-gang *et al.*, 2009; Singh, 1997). Due to its attractive shapes and colors, it is an ideal ornamental plant for gardens as well as cut flower (Barze-gar *et al.*, 2004). It has been known that growth regulators among the agriculture practices are the most favorable for promoting and improving plant growth (Eid and Abou-Leila, 2006). At present, due to their key role in the growth and developmental processes of plants, PGRs are used widely. Foliar sprays and drenches are the most common methods of application of growth regulators (Al-khassawneh *et al.*, 2006). Although drench method allows more uniform control of plant height and uses less active material and lower concentrations of the growth regulator, it has higher labor costs than foliar sprays (Al-khassawneh *et al.*, 2006).

Gibberellins (GAs) are a family of plant hormones controlling many aspects of plant growth and development, including stem elongation, germination, and the transition from vegetative growth to flowering (Thomas *et al.*, 2005; Gul *et al.*, 2006). Pre-sowing treatment of corms with GA₃ proved effective in increasing the flower number, plant height and corm yield of gladiolus (Bhattacharjee, 1984; Roychowdhury, 1989). It was reported that GA₃ significantly increased the length of flower stalk of *Rosa hybrid* cv. Poison (Hashemabadi and Zarchini, 2010). Gul *et al.* (2006) reported that GA₃ accelerated flowering and enhanced plant height of *Araucaria hetero-phylla*. Salicylic acid is considered as a potent plant hormone because of its diverse regulatory roles in plant metabolism (Popova *et al.*, 1997; Hayat and Ahmad, 2007). Exogenous application of SA may influence a range of diverse processes in plants, including seed germination, stomatal closure, ion uptake and transport, membrane permeability, photosynthetic and growth rate (El-Tayeb, 2005). Krizatsin (2-methyl phenoxy acetic acid) and mival (chlormethyl silatrane) are extensively used in Russia (Kazimirovskaya *et al.*, 2001). These compounds have hormone-like activity and their impact on the growth of cells, formation of embryonic organs and improvement of embryo growth has been demonstrated (Voronkov *et al.*, 2005; Voronkov *et al.*, 1978). It was found that soaking barley seeds in mival solution increased seed germination (Voronkov *et al.*, 2005). Furthermore, mival treatment significantly improves the yield of tomatoes (by 8–24%) and sugar beets (by 10%) (Voronkov and Baryshok, 2010). Krizatsin is the homolog of the herbicide 2, 4-D, but it has no toxicity effect on the plants. It is one of the main secondary metabolites such as vitamins A and E, and therefore, enhances the strength and stability of the cell membrane and also stimulates sexual activities in both plants and animals.

The main aim of the present study was to determine the effectiveness of GA₃, SA, KR and MI on vegetative and reproductive parameters of *Gladiolus hybridus*.

MATERIALS AND METHODS

The experiment was carried out in a greenhouse at the Faculty of Agriculture and Natural Resources, Persian Gulf University (PGU), Bushehr, Iran (29° 169' N, 51° 129' E, elevation, 65 m). The corms were disinfected with a benomyl fungicide (0.2%) for 10 minutes just before planting and then corms were divided into two groups. Half of corms were applied as a pre-plant drench for 8 h at 0, 25, 50 mg L⁻¹ for all chemical PGRs. The other half was used for spray treatments with SA, MI and KR at 0, 50, 100 mg L⁻¹ and GA₃ at 0, 25, 50 mg L⁻¹ (Nov. 15, 2008). The corms were planted on ridges (20 cm) at 8-10 cm depth. After harvesting flowering shoots in order to complete the growth of the corms, watering continued until leaves were yellow. Drip irrigation system was used for the experiment. After the appearance of floral stalk, traits including length of floral stalk, florets number, floret diameter, leaf area, leaf chlorophyll index and corm weight and volume were measured. Leaf area and leaf chlorophyll index were measured by portable leaf area meter (Model CI-202) and chlorophyll meter (SPAD), respectively. The experiment was planned as factorial design in a completely randomized design with four replicates. Treatments included PGRs solutions (GA₃, SA, MI and KR) at various concentrations with two methods of application

(drenched and foliar sprayed). All studied traits were subjected to analysis of variance. Statistical analysis of data was carried out by SAS and mean comparison was done by Duncan's multiple range test ($P < 0.05$).

RESULTS

Length of floral stalk

Data presented in Table 1 showed that the application methods and PGRs had no significant effect on the length of floral stalk. However, the interaction of application methods \times PGRs type \times concentration was significant ($P < 0.01$). The application of 50 mg L⁻¹ of GA₃ and SA as spray significantly increased the length of floral stalk of gladiolus by 105.37 and 105.25 cm as compared to control (94.75 cm), respectively (Fig. 1).

Floret diameter

The floret diameter was significantly ($P < 0.05$) affected by the interaction of application methods \times PGRs type \times concentration (Table 1). The highest floret diameter (14.25 mm) was obtained when the plants were sprayed with MI at 100 mg L⁻¹ (Fig. 2).

Florets number

The number of florets was significantly affected by the interaction of PGRs type \times application methods (Table 1). The maximum number of florets (16.25) was observed in the plants treated with MI at intermediate concentrations in both application methods (Fig. 3).

Table 1. Analysis of variance for *Gladiolus hybridus* to evaluate the effects of GA₃, SA, MI and KR treatments on studied traits.

S. O. V.	df	MS						
		Length of floral stalk	Floret diameter	Floret number	Leaf area	Corm weight	Corm volume	Leaf chlorophyll index
Application methods (A)	1	47.91 ^{ns}	0.041 ^{ns}	2.04 ^{ns}	311.3 ^{ns}	153.41**	38.64**	52.06 [*]
PGRs (B)	3	21.58 ^{ns}	1.52 ^{ns}	7.52**	435 [*]	1202.2**	807.55**	21.12 ^{ns}
Concentration (C)	2	199**	0.322 ^{ns}	6.50**	171.03 ^{ns}	1640.1**	2614.87**	36.81 ^{ns}
A \times B	3	42.70 ^{ns}	2.06 ^{ns}	1.40 ^{ns}	304.4 ^{ns}	285.6**	526.09**	45.70 [*]
A \times C	2	14.99 ^{ns}	0.01 ^{ns}	0.54 ^{ns}	82.55 ^{ns}	742.60**	512.46**	15.23 ^{ns}
B \times C	6	65.75**	1.05 ^{ns}	4.73**	826.2**	680.42**	509.34**	31.07 [*]
A \times B \times C	6	117.74**	1.91 [*]	1.52 ^{ns}	223.8 ^{ns}	119.97**	149.07**	24.94 ^{ns}
Error	72	17.85	0.784	1.27	136.07	2.72	2.56	11.95

*: Significant at $P < 0.05$; **: Significant at $P < 0.01$; ns: not significant.




Fig. 1. Interaction of application method \times PGRs type \times concentration for the length of floral stalk (D: drenched plant, S: sprayed plant).




Fig. 2. Interaction of PGRs × application methods × concentration for floret diameter.




Fig.3. Interaction of hormone × concentrations for floret number.

Leaf area

Leaf area of gladiolus cut flowers was significantly ($P < 0.05$) affected by PGRs (Table 1). In addition, leaf area was significantly ($P < 0.01$) affected by the interaction of PGRs type × concentration. The highest and the lowest leaf areas were related to the high concentrations of GA₃ and SA, respectively (Fig. 4).

Corm weight and volume

It was also revealed that application methods and different concentrations of PGRs had significant impact on the weight and volume of corm (Table 1). Moreover, these traits were significantly ($P < 0.01$) affected by the interaction of application methods × PGRs type × concentration. Plants treated with MI at 25 mg L⁻¹ had significantly higher weight and volume of corm.




Fig.4. Interaction of PGRs type × concentrations for leaf area.




Fig.5. Interaction of PGRs × application method for leaf chlorophyll index.

Leaf chlorophyll index

There was no significant difference among PGRs types in terms of leaf chlorophyll index (Table 1). However, leaf chlorophyll index was significantly ($P < 0.05$) affected by the interaction of PGRs type × concentration and also application methods × PGRs type (Table 1). The highest leaf chlorophyll index was observed in KR, SA and MI as drench application (Fig. 5).

DISCUSSION

In this study, it was found that PGRs treatments significantly improved some traits of gladiolus cut flowers as compared to control. Plants sprayed with GA₃ and SA at 50 mgL⁻¹ exhibited significantly longer floral stalk. Our findings are in agreement with Hassanpour Asil *et al.* (2011) on *Polianthes tuberosa* and Al-Khassawneh *et al.* (2006) on *Iris nigricans* who indicated drenching with GA₃ increased the floral stalk height as compared to the control. It may be attributed to the effect of GA₃ in stimulating and accelerating cell division, increasing cell elongation and enlargement, or both (Hartmann *et al.*, 1990). In contrast to our results, Hashemabadi and Zarchini (2010) on cut rose (*Rosa hybrida* cv. Poison) reported that SA did not show any significant effect on flower stem height as compared to the control. In this study, the highest and lowest leaf areas were achieved in plants treated with high GA₃ and SA doses, respectively. Our findings are in agreement with Emongor (2004) who reported that GA₃ application increased petiole length and leaf area. Also, consistent with our results, Hassanpour Asil *et al.* (2011) indicated that the application of 50 and 100 mg L⁻¹ of GA₃ significantly increased leaf length of *P. tuberosa* by 21.94 and 21.13 cm, respectively. Our results showed that plants sprayed with 100 mg L⁻¹ MI exhibited significantly higher floret diameter. It seems that MI (chloromethyl silatrane) increases floret diameter through stimulating the growth of cells (Voronkov *et al.*, 2005).

The maximum number of florets was obtained in the plants treated with intermediate concentrations of MI and GA₃ and high concentration of SA. Our findings agree with Muchopadhyay and Banker (1983) who mentioned that the GA₃ spray increased spike length and florets number per spike of *Polianthes*. In contrast to our results, Hassanpour Asil *et al.* (2011) reported that GA₃ reduced the number of florets.

In this research, the highest corm weight and volume was obtained when 25 mg L⁻¹ MI was applied as drench to cut flowers, whereas the application of GA₃ reduced these traits in both application methods. In contrast to our results, Ram *et al.*, (2002) reported that GA₃ increased corm weight of *Gladiolus*. This effectiveness of MI may be due to an increase in assimilate mobilization, as the water potential of the corm (the competitive sink) remained high aiding its growth and assimilate attraction.

CONCLUSION

Based on these results, it is concluded that the use of MI had major positive effects on florets number, floret diameter and corm weight and volume. Additionally, the application method showed no significant effect on the majority of traits in this experiment. SA showed the least effect among plant growth regulators applied to gladiolus cut flowers.

Literature Cited

- Al-khassawneh, N.M., Karam, N.S. and Shibli, R.A. 2006. Growth and flowering of black iris (*Iris nigricans* Dinsm.) flowering treatment with plant growth regulators. *Scientia Horticulturae*. 107:187-193.
- Barzegar, T., Khalighi, A. and Naderi, R. 2004. Effect of nitrogen rate and planting time on vegetative traits and corm yield of gladiolous (*Gladiolus grandiflora* L. cv. Oscar). *Seed and Plant Improvement Journal*. 22 (1): 55-65.
- Bhattacharjee, S.K. 1984. The effect of growth regulating chemicals on gladiolous. *Garten bauwissenschaften*. 49: 103-106.
- Eid, R.A. and Abou-Leila, B.H. 2006. Response of croton plants to gibberellic acid, benzyl adenine and ascorbic acid application. *World Journal of Agricultural Sciences*. 2:174-179.
- El-Tayeb, M.A. 2005. Response of barley grains to the interactive effect of salinity and salicylic acid. *Plant Growth Regulation*. 45:215–224.
- Emongor, V. E. 2004. Effect of gibberellic acid on post-harvest quality and vase life of gerbera cut flowers. *Journal of Agronomy*, 3: 191-195.
- Gul, H., Khattak, A.M. and Amin, N. 2006. Accelerating the growth of *araucaria heterophylla* seedlings through different gibberellic acid concentrations and nitrogen levels. *Journal of Agricultural and Biological Science*.1(2): 25-29.
- Hartmann, H.T., Kester, D.E. and Davies, F.T. 1990. *Plant propagation, principles and practices*. 5th ed. Prentice-Hall, Englewood Cliffs, NJ.
- Hassanpour Asil, M., Roein, Z. and Abbasi, J. 2011. Response of tuberose (*Polianthes tuberosa* L.) to gibberellic acid and benzyladenine. *Horticulture, Environment and Biotechnology*, 52(1):46-51.
- Hashemabadi, D. and Zarchini, M. 2010. Yield and quality management of rose (*Rosa hybrida* cv. Poison) with plant growth regulators. *Plant Omics Journal*. 3(6):167-171.
- Hayat, S. and Ahmad, A. 2007. *Salicylic acid - A Plant Hormone*. Springer. 401 p.
- Ji-gang, B., Pei-lei, X., Cheng-shun, Z. and Cai-yun, W. 2009. Effects of exogenous calcium on some postharvest characteristics of cut gladiolus. *Agricultural Sciences in China*. 8 (3): 293-303.
- Kazimirovskaya, V.B., D'yakov, V.M. and Voronkov, M.G. 2001. 1-(Chloromethyl) silatrane: a regulator of the functional activity of connective tissues. *Pharmaceutical Chemistry Journal*. 35 (9): 465-467.
- Muchopadhyay, A. and Banker, G.J. 1983. Regulation of growth and flowering in *Polianthes tuberosa* L. with gibberellic acid on ethrel spray. *Horticulture Science*.19: 149-152.
- Popova, L., Pancheva, T. and Uzunova, A. 1997. Salicylic acid: properties, biosynthesis and physiological role. *Bulgarian Journal of Plant Physiology*. 23: 85–93.
- Ram, R., Mukherjee, D. and Manuja, S.2002. Plant growth regulators affect the development of both corms and cormels in gladiolus. *Hort Science*, 37: 343-344.
- Roychowdhury, N. 1989. Effect of plant spacing and growth regulators on growth and flower yield of gladiolus grown under polythene tunnel. *Acta Horticulturae*, 246: 259-263.
- Singh, K.P. 1997. Improved agro-techniques for gladiolus. A review of work done in India. *Agricultural Reviews*. 18:212–238.
- Thomas, S.G., Rieu, I. and Steber, C.M. 2005. Gibberellin metabolism and signaling. *Vitamins and Hormones*. 72:289–338.
- Voronkov, M. G. and Baryshok, V. P. 2010. Atranes as a new generation of biologically active substances. *Herald of the Russian Academy of Sciences*. 80(6): 514-521.
- Voronkov, M. G., Dolmaa G., Tserenpil, Sh., Ugtakhbayar, O. and Chimidtsogzol, A. 2005. Stimulation of barley seed germination by micromolar aqueous solutions of silatrane and cresacin. *Doklady Biological Sciences* .404 (4): 562.
- Voronkov, M. G., Platonova, R. N., Svarinskaia, R., Karpova, N. I. and Diakov, V. 1978. Cytophysiological analysis of action of 1-(chloromethyl)-silatrane on meristem cells of higher-plants. *Doklady Akademii Nauk SSSR*, 242(6):1407-1410.